arXiv:1504.04195v2 [math.CO] 7 Jan 2016

Spectral conditions for Hamiltonicity of claw-free graphs

Binlong Li[a], Bo Ning[b]*and Xing Peng[b]

[a] Department of Applied Mathematics, Northwestern Polytechnical University,
Xi'an, Shaanxi 710072, P.R. China
[b] Center for Applied Mathematics, Tianjin University,
Tianjin, 300072, P.R. China
`libinlong@mail.nwpu.edu.cn; ningbo-maths@163.com; x2peng@tju.edu.cn`

Abstract

Fiedler and Nikiforov gave sufficient conditions for the existence of Hamilton paths and cycles in terms of spectral radii of a graph and its complement, which stimulated much later works on this topic. The class of claw-free graphs, which contains line graphs as a subclass, is an important type of graphs. Motivated by Fiedler and Nikiforov's works, in this paper we prove tight sufficient spectral conditions for Hamilton cycles in 2-connected claw-free graphs and Hamilton paths in connected claw-free graphs. Our strategy is to deduce spectral results from structural results, and main tools include Ryjácek's claw-free closure theory a characterized theorem of minimal 2-connected non-hamiltonian claw-free graphs due to Brousek, together with some spectral inequalities.

Keywords: Spectral radius; Q-index; Hamiltonicity; claw-free graph

Mathematics Subject Classification (2010): 05C50; 05C45; 05C35

1 Introduction

We will continue our recent works [15]. The main goal of this note is to give spectral conditions for Hamiltonicity of 2-connected claw-free graphs. For terminology and notation not defined but used, we refer the reader to [2, 5]. For a graph G, we denote by $e(G)$ the edge number of G and $\omega(G)$ the clique number of G. For two graphs G_1 and G_2, let $G_1 + G_2$ and $G_1 \vee G_2$ denote the *disjoint union* and *join* of G_1 and G_2, respectively.

Let G be a graph, A be the *adjacency matrix* of G, and D be the *degree matrix* of G. We call $Q = D + A$ the *signless Laplace matrix* of G. The *spectral radius* of G, denoted by $\mu(G)$, is the largest value among all eigenvalues of A; and the *signless Laplacian spectral radius* (or *Q-index*) of G, denoted by $q(G)$, is the largest value among all eigenvalues of Q.

A graph G is called *Hamiltonian* (*traceable*) if there is a Hamilton cycle (Hamilton path), i.e., a cycle (path) containing all vertices in G. The topic of finding spectral

---

[*] Corresponding author.

conditions for the existence of some kinds of subgraph in a graph has been well studied. In particular, sufficient spectral conditions for traceability and Hamiltonicity of graphs have received much attentions from graph theorists.

Fielder and Nikiforov [8] gave tight sufficient conditions for Hamilton paths and cycles in terms of spectral radii of a graph and its complement.

Theorem 1.1 (Fiedler and Nikiforov [8])**.** *Let G be a graph on n vertices.*
(1) If $\mu(G) \geq n-2$, then G is traceable unless $G = K_{n-1} + K_1$.
(2) If $\mu(G) > n-2$, then G is Hamiltonian unless $G = K_1 \vee (K_1 + K_{n-2})$.

Theorem 1.2 (Fiedler and Nikiforov [8])**.** *Let G be a graph on n vertices.*
(1) If $\mu(\overline{G}) \leq \sqrt{n-1}$, then G is traceable unless $G = K_{n-1} + K_1$.
(2) If $\mu(\overline{G}) \leq \sqrt{n-2}$, then G is Hamiltonian unless $G = K_1 \vee (K_1 + K_{n-2})$.

For Q-index, Zhou [18] and Liu et al. [11] gave some sufficient conditions for Hamilton cycles or Hamilton paths in terms of Q-indices of a graph and its complement. Since the connectedness is a necessary condition for a graph to be traceable, it is natural to consider the traceability of graphs in the class of connected graphs. Lu et al. [12] obtained spectral radius conditions for Hamilton paths of connected graphs. Ning and Ge [14] refined Lu et al.'s results.

Some works also focus on Hamiltonicity of some special classes of graphs. Date back to 1986, Simić and Kocić [17] determined maximal spectral radius among bicyclic Hamiltonian graphs with given number of edges. Bell and Rowlinson [1] studied similar problem for tricyclic Hamiltonian graphs. Zhou et al. [19] gave some spectral radius conditions for Hamiltonicity of planar graphs and outer-planar graphs. Lu et al. [12] and Liu et al. [11] prove spectral conditions for Hamilton paths and cycles in bipartite graphs.

A graph is *claw-free* if it contains no copy of claw ($K_{1,3}$) as an induced subgraph. Claw-free graphs, including line graphs, is an important type of graphs. There are lots of results on the Hamiltonian properties of claw-free graphs. The following is a well-known example, where N is the graph obtained from a triangle by adding three disjoint pendent edges.

Theorem 1.3 (Duffus et al [6])**.** *Let G be a claw-free graph with no induced copies of N.*
(1) If G is connected, then G is traceable.
(2) If G is 2-connected, then G is Hamiltonian.



Fig. 1. Graph EN_n.

In this paper, we use EN_n ($n \geq 6$) to denote the graph obtained from a complete graph K_{n-3} by adding three disjoint pendent edges, see Fig. 1. Note that $N = EN_6$. Recently, we proved some spectral conditions on traceability of connected claw-free graphs [15].

Theorem 1.4 (Ning and Li [15])**.** *Let G be a connected claw-free graph on n vertices. If $\mu(G) \geq n-4$, then G is traceable unless $G = EN_n$.*

Theorem 1.5 (Ning and Li [15])**.** *Let G be a connected claw-free graph on $n \geq 24$ vertices. If $\mu(\overline{G}) \leq \mu(\overline{EN_n})$, then G is traceable unless $G = EN_n$.*

For the Q-index, we have a similar result as follows:

Theorem 1.6. *Let G be a connected claw-free graph on $n \geq 18$ vertices. If $q(G) \geq q(EN_n)$, then G is traceable unless $G = EN_n$.*

The proof of Theorem 1.6 is postponed to Section 3.

Now we study the Hamiltonicity of 2-connected claw-free graphs under spectral conditions.

Before giving our main results, we first introduce some notations by Brousek. Following [4], we use $\mathcal{P}$ to denote the class of graphs obtained by two triangles $a_1a_2a_3a_1$ and $b_1b_2b_3b_1$, such that each pair $\{a_i, b_i\}$ is connected by a triangle or a path of order $k_i \geq 3$. We use P_{x_i,x_2,x_3} to denote the graph from $\mathcal{P}$, where $x_i = T$ if $\{a_i, b_i\}$ is connected by a triangle; and $x_i = k_i$ if $\{a_i, b_i\}$ is connected by a path of order k_i.


$P_{T,T,T}$ $P_{3,T,T}$ $P_{3,3,T}$ $P_{3,3,3}$

Fig. 2. 2-connected claw-free non-Hamiltonian graphs of order 9.

Brousek [4] characterized the minimal 2-connected claw-free non-Hamiltonian graphs.

Theorem 1.7 (Brousek [4])**.** *Every 2-connected claw-free non-Hamiltonian graph contains an induced subgraph isomorphic to a graph in $\mathcal{P}$.*

From Theorem 1.7, we can see that the smallest 2-connected claw-free non-Hamiltonian graphs have order 9, and there are exactly four such graphs, namely $P_{T,T,T}$, $P_{3,T,T}$, $P_{3,3,T}$ and $P_{3,3,3}$, see Fig. 2. We use $\mathcal{EP}_n$ ($n \geq 9$) to denote the set of graphs which are obtained from $P_{T,T,T}$, $P_{3,T,T}$, $P_{3,3,T}$ or $P_{3,3,3}$ by replacing a triangle by K_{n-6}. we use EP_n and EP'_n to denote the two graphs in $\mathcal{EP}_n$ obtained from $P_{T,T,T}$ by replacing a triangle by K_{n-6}, see Fig. 3. Note that each graph in $\mathcal{EP}_n$ is a subgraph of EP_n or EP'_n.



EP_n EP'_n

Fig. 3. Graphs EP_n and EP'_n

Our main results are closely related to Brousek's characterization and given as follows. Let G be a 2-connected claw-free graph on n vertices:

Theorem 1.8. *If $n \geq 33$ and $\mu(G) \geq \mu(EP_n)$, then G is Hamiltonian unless $G = EP_n$.*

Theorem 1.9. *If $n \geq 51$ and $q(G) \geq q(EP_n)$, then G is Hamiltonian unless $G = EP_n$.*

Theorem 1.10. *If $n \geq 219$ and $\mu(\overline{G}) \leq \mu(\overline{EP'_n})$, then G is Hamiltonian unless $G = EP'_n$.*

The organization of this note is as follows. In Section 2, we will introduce useful tools from structural graph theory and spectral inequalities from spectral graph theory. In Section 3, we will give proofs of our main results.

2 Preliminaries

In this section, we mainly give useful tools from structural graph theory and spectral graph theory.

Generally speaking, most of existed researches on this topic appeared the following characterization: The spectral part can always be deduced from the non-spectral part, and the main technique is aid of the powerful and well-chosen closure theory (see [8, 12] for example). This is also our strategy here (together with some other structural and spectral tools).

Since we study claw-free graphs here, we will use the claw-free closure theory introduced by Ryjáček [16]. To ensure the completeness of our text, all necessary terminology and notations about claw-free closure theory are included. For other more information, see [16].

Let G be a claw-free graph. Following [16], for a vertex $x \in V(G)$, if the neighborhood of x induces a connected but not complete subgraph of G, then x is called *eligible* in G. Set $B_G(x) = \{uv : u, v \in N(x), uv \notin E(G)\}$. The graph G'_x, constructed by $V(G'_x) = V(G)$ and $E(G'_x) = E(G) \cup B_G(x)$, is called the *local completion of G at x*. The *closure* of G, denoted by $cl(G)$, is defined by a sequence of graphs $G_1, G_2, \ldots, G_t$, and vertices $x_1, x_2 \ldots, x_{t-1}$ such that
(1) $G_1 = G$, $G_t = cl(G)$;

(2) x_i is an eligible vertex of G_i, $G_{i+1} = (G_i)'_{x_i}$, $1 \le i \le t-1$; and
(3) $cl(G)$ has no eligible vertices.
Ryjáček [16] proved that the closure of a claw-free graph is also claw-free.

Theorem 2.1 (Brandt et al. [3]). *Let G be a claw-free graph. Then G is traceable if and only if $cl(G)$ is traceable.*

Theorem 2.2 (Ryjáček [16]). *Let G be a claw-free graph. Then G is Hamiltonian if and only if $cl(G)$ is Hamiltonian.*

A claw-free graph G is said to be *closed* if $cl(G) = G$. It is not difficult to see that for every vertex x of a closed graph G, $N(x)$ is either a clique, or the disjoint union of two cliques in G (see [16]). We prove the following lemma on Hamiltonicity of closed claw-free graphs.

Lemma 1. *Let G be a 2-connected closed claw-free graph on n vertices. If there are two nonadjacent vertices $u, v \in V(G)$ such that $d(u) + d(v) \ge n$, then G is Hamiltonian.*

Proof. Since $uv \notin E(G)$ and $d(u)+d(v) \ge n$, u and v have at least two common neighbors. Note that any two common neighbors of u and v are nonadjacent; otherwise they will be eligible vertices of G. If u and v have three common neighbors, then there will be a claw with center u. This implies that u and v have exactly two common neighbors x, y, say, and moreover, every vertex in $V(G)\backslash\{u, v, x, y\}$ is adjacent either u or v. Note that $N(u)$ ($N(v)$) is the disjoint union of two cliques. This implies that G consists of four cliques C_1, C_2, C_3 and C_4 containing $\{u, x\}, \{x, v\}, \{v, y\}$ and $\{y, u\}$, respectively. It is easy to check that G has a Hamilton cycle. □

Lemma 2. *Let k be an integer and G be a closed claw-free non-Hamiltonian graph on $n \ge 2k$ vertices. If*

$$e(G) \ge \binom{n-k-1}{2} + \binom{k+2}{2} + 1,$$

then $\omega(G) \ge n - k$.

Proof. A vertex v is a *heavy vertex* of G if $d(v) \ge n/2$. By Lemma 1, any two heavy vertices are adjacent in G. Let C be a maximum clique of G containing all heavy vertices and let $H = G - C$. Set $t = |C|$.

Let v be an arbitrary vertex in H. Then v is nonadjacent to at least one vertex in C. If there are at least two neighbors of v in C, then there will be eligible vertices, contradicting the fact that G is closed. This implies that every vertex in H has at most one neighbor in C.

We first assume that $1 \le t \le (n+1)/2$. Recall that every vertex in H has degree at most $(n-1)/2$ in G. Hence

$$\sum_{v \in V(H)} (d_C(v) + d(v)) \le (n-t)\left(\frac{n-1}{2} + 1\right) = \frac{(n-t)(n+1)}{2}.$$

and

$$\begin{aligned}
e(G) &= e(G[C]) + \frac{\sum_{v\in V(H)}(d_C(v)+d(v))}{2}\\
&\leq \binom{t}{2} + \frac{(n-t)(n+1)}{4}\\
&= \frac{1}{2}t^2 - \frac{n+3}{4}t + \frac{n(n+1)}{4}\\
&\leq \frac{n^2-1}{4}\\
&\leq \binom{n-k-1}{2} + \binom{k+2}{2} < e(G),
\end{aligned}$$

a contradiction.

Now we assume that $n/2+1 \leq t \leq n-k-1$. Recall that every vertex in H has at most one neighbor in C. Thus

$$\begin{aligned}
e(G) &= e(G[C]) + e(H) + e(C, V(H))\\
&\leq \binom{t}{2} + \binom{n-t}{2} + (n-t)\\
&= \binom{t}{2} + \binom{n-t+1}{2}\\
&\leq \binom{n-k-1}{2} + \binom{k+2}{2} < e(G),
\end{aligned}$$

a contradiction.

This implies that $t \geq n-k$, and then $\omega(G) \geq t \geq n-k$. $\square$

Lemma 3. *Let G be a connected closed claw-free non-traceable graph on n vertices. If $\omega(G) \geq n-3$, then $G = EN_n$.*

Proof. Since G is connected claw-free and non-traceable, by Theorem 1.3, G contains an induced subgraph H isomorphic to N. Let C be a maximum clique of G. If $|C \cap V(H)| \leq 2$, then $|C \cup V(H)| \geq n-3+6-2 = n+1$, a contradiction. Hence $|C \cap V(H)| \geq 3$. Since $\omega(H) = 3$, $|C \cap V(H)| \leq 3$. Thus $|C \cap V(H)| = 3$ and C contains the (unique) triangle of H. Thus $EN_n \subseteq G$. Note that the graph obtained from EN_n by adding any edge is traceable. Thus $G = EN_n$. The proof is complete. $\square$

Lemma 4. *Let G be a connected claw-free graph on $n \geq 6$ vertices. If $e(G) \geq n(n-9)/2+21$, then G is traceable, unless $G \subseteq EN_n$.*

Proof. Let $G' = cl(G)$. If G' is traceable, then so is G by Theorem 2.1. Now we assume that G' is non-traceable. Clearly $e(G') \geq e(G) \geq n(n-9)/2+21$. Since G' is closed, by Lemma 2, $\omega(G) \geq n-3$; and by Lemma 3, $G' = EN_n$. Thus $G \subseteq EN_n$. $\square$

Lemma 5. *Let G be a 2-connected closed claw-free non-Hamiltonian graph on n vertices. If $\omega(G) \geq n-6$, then $G \in \mathcal{EP}_n$.*

Proof. Since G is 2-connected closed claw-free and non-Hamiltonian, by Theorem 1.7, G contains an induced subgraph H isomorphic to a graph in $\mathcal{P}$. Let C be a maximum clique of G. If $|C \cap V(H)| \leq 2$, then $|C \cup V(H)| \geq n-6+9-2 = n+1$, a contradiction. Hence $|C \cap V(H)| \geq 3$. Since $\omega(H) = 3$, $|C \cap V(H)| \leq 3$. Thus $|C \cap V(H)| = 3$. If $|V(H)| \geq 10$, then $|C \cup V(H)| \geq n-6+10-3 = n+1$, a contradiction. Hence $|V(H)| = 9$ and G is a supergraph of a graph in $\mathcal{EP}_n$. If $G \notin \mathcal{EP}_n$, then easily to check that G is Hamiltonian. This implies that $G \in \mathcal{EP}_n$. The proof is complete. $\square$

Lemma 6. *Let G be a connected claw-free graph on $n \geq 12$ vertices. If $e(G) \geq n(n-15)/2+57$, then G is traceable, unless $G \subseteq EP_n$ or EP'_n.*

Proof. Let $G' = cl(G)$. If G' is Hamiltonian, then so is G by Theorem 2.2. Now we assume that G' is non-Hamiltonian. Clearly $e(G') \geq e(G) \geq n(n-15)/2+57$. By Lemma 2, $\omega(G) \geq n-6$; and by Lemma 5, $G' \in \mathcal{EP}_n$. This implies that G is a subgraph of a graph in $\mathcal{EP}_n$. Note that every graph in $\mathcal{EP}_n$ is either a subgraph of EP_n or a subgraph of EP'_n. Thus $G \subseteq EP_n$ or EP'_n. $\square$

We also need several spectral inequalities as follows.

Theorem 2.3 (Hong et al. [10]). *Let G be a graph of order n without isolated vertices. Then*

$$\mu(G) \leq \sqrt{2e(G)-n+1}.$$

Theorem 2.4 (Feng and Yu [7]). *Let G be a graph on n vertices. Then*

$$q(G) \leq \frac{2e(G)}{n-1}+n-2.$$

Theorem 2.5 (Hofmeister [9]). *Let G be a graph. Then*

$$\mu(G) \geq \sqrt{\frac{\sum_{v\in V(G)} d^2(v)}{n}}.$$

The following fact can be calculated by matrix analysis, together with the aid of Maple.

Lemma 7. *For $n \geq 10$, we have*
(1) $\mu(EP_n) > \mu(EP'_n) > \mu(K_{n-6}) = n-7$;
(2) $q(EP_n) > q(EP'_n) > q(K_{n-6}) = 2n-14$;
(3) $\mu(\overline{EP'_n}) < \mu(\overline{EP_n}) < \mu(K_6 \vee (n-6)K_1) = (5+\sqrt{24n-119})/2$.

Proof. Since proofs of inequalities (1), (2), and (3) are quite similar, we will give details for the proof of inequality (1) here and sketch others.
Proof of inequality (1): Let μ_1 (resp. μ_2) be the largest eigenvalue of the adjacency matrix of EP_n (resp. EP'_n). We will compute the characteristic equation of their adjacency matrices directly. Let x (resp. y) be the eigenvector corresponding to μ_1 (resp. μ_2). We observe EP_n has only four types of vertices with respect to their degrees, namely vertices with degree two, vertices with degree four, vertices with degree $n-4$, and vertices with

degree $n-7$. Furthermore, if two vertices u and v have the same degree, then $x_u = x_v$ by the symmetry. Similar observations also hold for the graph EP'_n.

Let u be a degree two vertex in EP_n, v be a degree four vertex in EP_n, w be a degree $n-4$ vertex in EP_n, and z be a degree $n-7$ vertex in EP_n. If A is the adjacency matrix of EP_n, then we have $Ax = \mu_1 x$. In particular, we have $(Ax)_u = \mu_1 x_u$, $(Ax)_v = \mu_1 x_v$, $(Ax)_w = \mu_1 x_w$, and $(Ax)_z = \mu_1 x_z$. Recall the observation above. We get the following system of linear equations:

$$x_v + x_w = \mu_1 x_u \tag{1}$$

$$x_u + 2x_v + x_w = \mu_1 x_v \tag{2}$$

$$x_u + x_v + 2x_w + (n-9)x_z = \mu_1 x_w \tag{3}$$

$$3x_w + (n-10)x_z = \mu_1 x_z \tag{4}$$

Viewing x_w as the free variable and solving for x_v and x_u from (1) and (2), we get

$$x_v = \frac{\mu_1 + 1}{\mu_1^2 - 2\mu_1 - 1} x_w \tag{5}$$

$$x_u = \frac{\mu_1 - 1}{\mu_1^2 - 2\mu_1 - 1} x_w. \tag{6}$$

Solving for x_z from (4), we get

$$x_z = \frac{3}{\mu_1 - n + 10} x_w. \tag{7}$$

Putting (5), (6), and (7) in (3), we obtain

$$\frac{\mu_1 - 1}{\mu_1^2 - 2\mu_1 - 1} x_w + \frac{\mu_1 + 1}{\mu_1^2 - 2\mu_1 - 1} x_w + 2x_w + \frac{3(n-9)}{\mu_1 - n + 10} x_w = \mu_1 x_w.$$

The Perron-Frobenius theorem implies that all entries in x are positive, so we can cancel x_w from both sides of the equation above. Simplify the resulted equation. We get the characteristic equation

$$(\mu_1 - n + 10)(\mu_1^3 - 4\mu_1^2 + \mu_1 + 2) - (3n - 27)(\mu_1^2 - 2\mu_1 - 1) = 0 \tag{8}$$

For EP'_n, using the same idea, we get a new system of linear equations

$$2y_v = \mu_2 y_u \tag{9}$$

$$y_w + 2y_v + y_u = \mu_2 y_v \tag{10}$$

$$2y_v + y_w + (n-8)y_z = \mu_2 y_w \tag{11}$$

$$2y_w + (n-9)y_z = \mu_2 y_z \tag{12}$$

From (9), (10), and (11), we get

$$y_v = \frac{\mu_2}{2} y_u$$

$$y_w = (\frac{\mu_2^2}{2} - \mu_2 - 1) y_u \tag{13}$$

$$y_z = \frac{(\mu_2^2 - 2\mu_2 - 2)(\mu_2 - 1) - 2\mu_2}{2n - 16} y_u \tag{14}$$

(13) together with (12) tell us

$$y_z = \frac{\mu_2^2 - 2\mu_2 - 2}{\mu_2 - n + 9} y_u \tag{15}$$

Recall $y_u > 0$. Equalizing (14) and (15) followed by cancelling y_u, we get

$$(\mu_2 - n + 9)\big((\mu_2^2 - 2\mu_2 - 2)(\mu_2 - 1) - 2\mu_2\big) - (2n - 16)(\mu_2^2 - 2\mu_2 - 2) = 0 \tag{16}$$

Define functions

$$f(x) = (x - n + 10)(x^3 - 4x^2 + x + 2) - (3n - 27)(x^2 - 2x - 1) \tag{17}$$

$$g(x) = (x - n + 9)\big((x^2 - 2x - 2)(x - 1) - 2x\big) - (2n - 16)(x^2 - 2x - 2) \tag{18}$$

Let $s = n - 7 + \frac{4}{(n-7)^2}$ and $t = n - 7 + \frac{7}{(n-7)^2}$. Tedious calculus together with Maple program can confirm $f(t) < 0$ and $g(s) < 0 < g(t)$ for all $n \geq 12$. Since K_{n-6} is a subgraph of EP_n and EP'_n, we get $\mu_1, \mu_2 \geq n - 7$. By Theorem 2.3, we get $\mu_1, \mu_2 < n - 5$. One can observe the gap between the largest eigenvalue and the second largest eigenvalue of the adjacency matrix of both EP_n and EP'_n is at least 2. Therefore, we obtain for all $n \geq 12$

$$\mu_1 > t \text{ and } s < \mu_2 < t,$$

which proves (1) for $n \geq 12$. For $n = 10, 11$, we can check the inequality (1) directly using Maple program.

Proof of inequality (2): Let $f(x)$ (resp. $g(x)$) be the characteristic polynomial of the signless Laplace matrix of EP_n (resp. EP'_n). Using the same idea as proving the inequality (1) (details will be given in Appendix A), we get

$$f(x) = (x - 2n + 17)\big((x - n + 3)(x^2 - 8x + 11) - (2x - 6)\big) - (3n - 27)(x^2 - 8x + 11)$$

$$\begin{aligned} g(x) &= (x^2 - 8x + 12)\left((x - 2n + 16)(x - n + 4) - (2n - 16)\right) \\ &\quad - (2x - 4n + 32)(2x - n + 2) - 4n + 32 \end{aligned}$$

We choose $s = 2(n - 7) + \frac{4}{n}$ and $t = 2(n - 7) + \frac{6}{n-7}$. Using basic calculus (under the help of Maple program) can verify $f(t) < 0$ and $g(s) < 0 < g(t)$ for all $n \geq 27$, which proves the inequality (2) for $n \geq 27$. For $10 \leq n \leq 26$, we can confirm the inequality using Maple program directly.

Proof of inequality (3): We use $f(x)$ (resp. $g(x)$) to denote the characteristic polynomial of the adjacency matrix of $\overline{EP_n}$ (resp. $\overline{EP'_n}$). We can obtain the formula for $f(x)$ and $g(x)$ by the same argument as we did for proving inequality (1) (the details will be presented in Appendix B). We get the following

$$f(x) = x^3 - (2x^2 + 4x + 8) - (6n - 54)(x + 1)$$

$$g(x) = (x^2 - 4n + 32)(x^2 - 2) - (2x^2 + x)(x + 2) - (2n - 16)(x^2 + x + 2)$$

Let $s = \sqrt{n(n - 6)}$ and $t = \sqrt{n(n - 6)} + 1.3$. With the assistance of Maple program, we can show $f(t) < 0$ and $g(s) < 0 < g(t)$ for all $n \geq 55$ using calculus. We proved the inequality (3) for $n \geq 55$. For the case of $10 \leq n \leq 54$, we can verify the inequality (3) using Maple program straightforwardly. $\square$

3 Proofs of the theorems

Proof of Theorem 1.6. Note that $q(EN_n) \geq q(K_{n-3}) \geq 2n-8$. By Theorem 2.4,

$$2n-8 \leq q(G) \leq \frac{2e(G)}{n-1} + n - 2.$$

One can get that

$$e(G) \geq \frac{(n-1)(n-6)}{2} \geq \frac{n(n-9)}{2} + 21$$

($n \geq 18$). By Lemma 4, G is traceable or $G \subseteq EN_n$. But if $G \subset EN_n$, then $q(G) < q(EN_n)$, a contradiction. Thus $G = EN_n$. The proof is complete.

Proof of Theorem 1.8. By Lemma 7 and Theorem 2.3,

$$n-7 \leq \mu(G) \leq \sqrt{2e(G)-n+1}.$$

One can get that

$$e(G) \geq \frac{(n-7)^2+n-1}{2} \geq \frac{n(n-15)}{2} + 57$$

($n \geq 33$). By Lemma 6, G is Hamiltonian or $G \subseteq EP_n$ or EP'_n. But if $G \subset EP_n$ or $G \subseteq EP'_n$, then $\mu(G) < \mu(EP_n)$, a contradiction. Thus $G = EP_n$. The proof is complete.

Proof of Theorem 1.9. By Lemma 7 and Theorem 2.4,

$$2n-14 \leq q(G) \leq \frac{2e(G)}{n-1} + n - 2.$$

One can get that

$$e(G) \geq \frac{(n-1)(n-12)}{2} \geq \frac{n(n-15)}{2} + 57$$

($n \geq 51$). By Lemma 6, G is Hamiltonian or $G \subseteq EP_n$ or EP'_n. But if $G \subset EP_n$ or $G \subseteq EP'_n$, then $q(G) < q(EP_n)$, a contradiction. Thus $G = EP_n$. The proof is complete.

Proof of Theorem 1.10. Let $G' = cl(G)$. If G' is Hamiltonian, then so is G by Theorem 2.2. Now we assume that G' is not Hamiltonian. By Lemma 1, every two nonadjacent vertices u, v have degree sum at most $n-1$, i.e.,

$$d_{\overline{G'}}(u) + d_{\overline{G'}}(v) \geq 2(n-1) - (n-1) = n-1.$$

By Theorem 2.5,

$$\mu(\overline{G}) \geq \mu(\overline{G'}) \geq \sqrt{\frac{\sum_{v\in V(G)} d^2_{\overline{G'}}(v)}{n}} = \sqrt{\frac{\sum_{uv\in E(\overline{G'})}(d_{\overline{G'}}(u)+d_{\overline{G'}}(v))}{n}} \geq \sqrt{\frac{(n-1)e(\overline{G'})}{n}}.$$

By Lemma 7,

$$\frac{5+\sqrt{24n-119}}{2} \geq \mu(\overline{G}) \geq \sqrt{\frac{(n-1)e(\overline{G'})}{n}}.$$

One can get that

$$e(G') = \binom{n}{2} - e(\overline{G'}) \geq \binom{n}{2} - \left(\frac{5+\sqrt{24n-119}}{2}\right)^2 \cdot \frac{n}{n-1} > \frac{n(n-15)}{2} + 56$$

($n \geq 219$). By Lemma 6, $G \subseteq G' \subseteq EP_n$ or EP'_n. But if $G \subseteq EP_n$ or $G \subset EP'_n$, then $\mu(\overline{G}) < \mu(EP'_n)$, a contradiction. Thus $G = EP'_n$. The proof is complete. □

References

[1] F.K. Bell, P. Rowlinson, On the index of tricycle Hamiltonian graphs, Proceedings of the Edinburgh Mathematical Society 33 (1990) 233–240.

[2] J.A. Bondy, U.S.R. Murty, Graph theory, Graduate Texts in Mathematics 244, Springer, New York, 2008.

[3] S. Brandt, O. Favaron, Z. Ryjáček, Closure and stable hamiltonian properties in claw-free graphs, J. Graph Theory 34 (1) (2000) 30–41.

[4] J. Brousek, Minimal 2-connected non-hamiltonian claw-free graphs, Discrete Math. 191 (1998) 57–64.

[5] D. Cvetkovi, M. Doob, H. Sachs, Spectra of Graphs, VEB Deutscher Verlag der Wissenschaften, Berlin, 1980.

[6] D. Duffus, M. Jacboson, R.J. Gould, Forbidden subgraphs and the hamiltonian theme, in: The Theory and Applications of Graphs (Wiley, New York 1981) 297–316.

[7] L. Feng, G. Yu. On three conjectures involving the signless Laplacian spectral radius of graphs, Publications de l'Institut Mathématique 85 (99) (2009) 35–38.

[8] M. Fiedler, V. Nikiforov, Spectral radius and Hamiltonicity of graphs, Linear Algebra Appl. 432 (9) (2010) 2170–2173.

[9] M. Hofmeisfer, Spectral radius and degree sequence, Mathematische Nachrichten 139 (1988) 37–44.

[10] Y. Hong, J.L Shu, K. Fang, A sharp upper bound of the spectral radius of graphs, J. Combin. Theory Ser. B 81 (2001) 177–183.

[11] R. Liu, W.C. Shiu, J. Xue, Sufficient spectral conditions on Hamiltonian and traceable graphs, Linear Algebra Appl. 467 (2015) 254–266.

[12] M. Lu, H. Liu, F. Tian, Spectral radius and Hamiltonian graphs, Linear Algebra Appl. 437 (7) (2012) 1670–1674.

[13] V. Nikiforov, Some inequalities for the largest eigenvalue of a graph, Combin. Probab. Comput. 11 (2002) 179–189.

[14] B. Ning, J. Ge, Spectral radius and Hamiltonian properties of graphs, Linear Multilinear Algebra 63 (8) (2015) 1520–1530.

[15] B. Ning, B. Li, Spectral radius and traceability of connected claw-free graphs, Filomat, Published Online.

[16] Z. Ryjáček, On a closure concept in claw-free graphs, J. Combin. Theory Ser. B 70 (1997) 217–224.

[17] S.K. Simić, V.Lj. Kocić, On the largest eigenvalue of some homeomorphic graphs, Publications de l'Institut Mathématique 40 (54) (1986) 3–9.

[18] B. Zhou, Signless Laplacian spectral radius and Hamiltonicity, Linear Algebra Appl. 432 (2/3) (2010) 566–570.

[19] J. Zhou, C. Lin, G. Hu, Spectral radius of Hamiltonian planar graphs and outerplanar graphs, Tsinghua Science and Technology 6 (4) (2001) 350–354.

Appendix A: Let q_1 (resp. q_2) be the largest eigenvalue of the signless Laplacian matrix of EP_n (resp. EP'_n). Let x (resp. y) be the eigenvector corresponding to q_1 (resp. q_2). Pick u as a degree two vertex in EP_n, v as a degree four vertex in EP_n, w as a degree $n-4$ vertex in EP_n, and z as a degree $n-7$ vertex in EP_n. If Q is the adjacency matrix of EP_n, then we have $Qx = q_1x$. Recall the symmetry of vertices with the same degree. We get the following system of linear equations.

$$2x_u + x_v + x_w = q_1x_u \tag{19}$$

$$6x_v + x_u + x_w = q_1x_v \tag{20}$$

$$(n-3)x_w + x_u + x_v + (n-9)x_z = q_1x_w \tag{21}$$

$$(2n-17)x_z + 3x_w = q_1x_z \tag{22}$$

We view x_w as a variable and solve for x_u, x_v, and x_z from (19), (20),(22). We get

$$x_u = \frac{q_1-5}{q_1^2-8q_1+11}x_w \tag{23}$$

$$x_v = \frac{q_1-1}{q_1^2-8q_1+11}x_w \tag{24}$$

$$x_z = \frac{3}{q_1-2n+17}x_w \tag{25}$$

Putting (23), (24), and (25) in (21), we get

$$\frac{q_1-5}{q_1^2-8q_1+11}x_w + \frac{q_1-1}{q_1^2-8q_1+11}x_w + \frac{3n-27}{q_1-2n+17}x_w = (q_1-n+3)x_w.$$

Cancel x_w from both sides of the equation above and simplify it. We get q_1 must satisfy the following equation

$$(3n-27)(q_1^2-8q_1+11) = (q_1-2n+17)((q_1-n+3)(q_1^2-8q_1+11)-(2q_1-6)),$$

which gives the characteristic polynomial of q_1.

For EP'_n, we select vertices u, v, w and z similarly. If Q is the signless Laplacian matrix of EP'_n, then we have $Qy = q_2y$. We obtain the following system of inequalities:

$$2y_u + 2y_v = q_2y_u \tag{26}$$

$$6y_v + y_u + y_w = q_2y_v \tag{27}$$

$$(n-4)y_w + 2y_v + (n-8)y_z = q_2y_w \tag{28}$$

$$(2n-16)y_z + 2y_w = q_2y_z \tag{29}$$

Using y_u to express y_v, y_w, and y_z in (26), (27), and (28), we get

$$y_v = \frac{q_2 - 2}{2} y_u \tag{30}$$

$$y_w = \left(\frac{1}{2}(q_2 - 2)(q_2 - 6) - 1\right) y_u \tag{31}$$

$$y_z = \frac{1}{n-8}\left((q_2 - n + 4)\left(\frac{1}{2}(q_2 - 2)(q_2 - 6) - 1\right) - (q_2 - 2)\right) y_u \tag{32}$$

We bring 30, 31, and 32 in 29 followed by cancelling y_u and simplification. We get q_2 must satisfy the following equation

$$(q_2^2 - 8q_2 + 12)\left((q_2 - 2n + 16)(q_2 - n + 4) - (2n - 16)\right) - (2q_2 - 4n + 32)(2q_2 - n + 2) - 4n + 32 = 0$$

which gives the characteristic polynomial of the signless Laplacian matrix of EP'_n.

Appendix B: For $\overline{EP_n}$, we let v be a vertex with degree $n-5$, u be a vertex with degree $n-3$, w be a vertex with degree 4, and z be a vertex with degree 6. We use μ_1 to denote the largest eigenvalue of the adjacency matrix A of EP'_n. If x is the eigenvector of μ_1, then we have $Ax = \mu_1 x$. Recall the symmetry between vertices with the same degree. We get

$$3x_u + 3x_v = \mu_1 x_z \tag{33}$$

$$2x_u + 2x_v = \mu_1 x_w \tag{34}$$

$$2x_u + 2x_w + (n-9)x_z = \mu_1 x_v \tag{35}$$

$$2x_u + 2x_v + 2x_w + (n-9)x_z = \mu_1 x_u \tag{36}$$

(33)-(34) gives

$$x_u + x_v = \mu_1 x_z - \mu_1 x_w \tag{37}$$

(36)-(35) yields

$$2x_v = \mu_1 x_u - \mu_1 x_v \tag{38}$$

We form a new system of linear equations using (34), (37), and (38). We solve for x_u, x_w, and x_z from the new system and we get

$$x_u = \frac{\mu_1 + 2}{\mu_1} x_v$$
$$x_w = \left(\frac{2(\mu_1 + 2)}{\mu_1^2} + \frac{2}{\mu_1}\right) x_v$$
$$x_z = \left(\frac{3(\mu_1 + 2)}{\mu_1^2} + \frac{3}{\mu_1}\right) x_v$$

We put expressions of x_u, x_w, and x_z in (35). Then we cancel x_v from both sides of the resulted equation and simplify it. We get

$$\mu_1^3 - (2\mu_1^2 + 4\mu_1 + 8) - (6n - 54)(\mu_1 + 1) = 0$$

which completes the proof.

For $\overline{EP_n'}$, we use the same assumptions as $\overline{EP_n}$. Let μ_2 be the largest eigenvalue and y be the corresponding eigenvector. We have the following system of linear equations.

$$4y_v + 2y_u = \mu_2 y_z \tag{39}$$

$$2y_v + 2y_u = \mu_2 y_w \tag{40}$$

$$y_u + y_v + y_w + (n-8)y_z = \mu_2 y_v \tag{41}$$

$$y_u + 2y_v + 2y_w + (n-8)y_z = \mu_2 y_u \tag{42}$$

(42)-(41) gives

$$y_v + y_w = \mu_2 y_u - \mu_2 y_v \tag{43}$$

(39)-(40) gives

$$2y_v = \mu_2 y_z - \mu_2 y_w \tag{44}$$

We solve for y_u, y_w, and y_z from the system consisting of equations (40), (43), and (44). We obtain

$$\begin{aligned} y_u &= \frac{\mu_2^2 + \mu + 2}{\mu_2^2 - 2} y_v \\ y_w &= \left(\frac{2}{\mu_2} + \frac{2(\mu_2^2 + \mu + 2)}{\mu_2(\mu_2^2 - 2)} \right) y_v \\ y_z &= \left(\frac{4}{\mu_2} + \frac{2(\mu_2^2 + \mu + 2)}{\mu_2(\mu_2^2 - 2)} \right) y_v \end{aligned}$$

We bring expressions of y_u, y_w, and y_z in (41), cancel y_v, and simplify the equation. We get μ_2 must satisfy the following equation

$$(\mu_2^2 - 4n + 32)(\mu_2^2 - 2) - (2\mu_2^2 + \mu_2)(\mu_2 + 2) - (2n - 16)(\mu_2^2 + \mu_2 + 2) = 0,$$

which completes the proof.